arXiv:1212.2337v1 [math.CV] 11 Dec 2012

The Hele-Shaw Flow and Moduli of Holomorphic Discs

Julius Ross and David Witt Nyström

December 12, 2012

Abstract

We present a new connection between the Hele-Shaw flow, also known as two-dimensional (2D) Laplacian growth, and the theory of holomorphic discs with boundary contained in a totally real submanifold. Using this we prove short time existence and uniqueness of the Hele-Shaw flow with varying permeability both when starting from a smooth Jordan domain and when starting from a point. Applying the same ideas we prove that the moduli space of smooth quadrature domains is a smooth manifold whose dimension we also calculate, and we give a local existence theorem for the inverse potential problem in the plane.

1 Introduction

The Hele-Shaw flow is a model for describing the propagation of fluid in a Hele-Shaw cell. Such a cell consists of two parallel plates separated by a small gap, and the fluid is confined to the narrow space between them. This moving boundary model has been intensely studied for over a century, and is a paradigm for understanding more complicated systems such as the flow of water in a porous media, the melting of ice and models of tumour growth.



Figure 1: Cross section of a Hele-Shaw cell

This flow is essentially two-dimensional, and one identifies the region occupied by fluid with a subset of the plane $\mathbb{R}^2$. In this paper we will consider the case of a viscous incompressible fluid that is surrounded by air while new fluid is injected at a constant rate at the origin. If p is the pressure in the fluid then the velocity V of the fluid at that point (or really the mean velocity along the gap between the plates) is calculated as

$$V = -\nabla p. \tag{1}$$

Here we have neglected some physical constants. The pressure in the fluid is harmonic except at the origin, where it will have a logarithmic singularity, and is zero on the boundary since we assume that both the air pressure and surface tension is zero.

To model this, let $\{\Omega_t\}$ be a family of domains in $\mathbb{R}^2$ containing the origin, and let p_t denote the function which is zero on $\partial\Omega_t$ and which solves

$$\Delta p_t = -\delta_0$$

on Ω_t, where δ_0 is the Dirac measure at the origin. Then Ω_t is called a solution to the (classical) Hele-Shaw flow if for all t

$$V_t = -\nabla p_t \quad \text{on } \partial\Omega_t, \tag{2}$$

where V_t is the normal outward velocity of $\partial\Omega_t$. Clearly we need some regularity of the family $\{\Omega_t\}$ in order for this to make sense, thus the question of local existence of a classical solution to the Hele-Shaw flow with a given initial domain Ω_0 is non-trivial.



Figure 2: Dynamics of the Hele-Shaw flow

Similarly one can model the case of a Hele-Shaw cell where the fluid moves through a porous material. If the permeability of this material is given by a positive function κ then, by Darcy's law, the equation of motion of a fluid in the cell becomes

$$V_t = -\kappa\nabla p_t \quad \text{on } \partial\Omega_t, \tag{3}$$

with p_t defined as before. In [25] this situation is considered, where it goes under the name of elliptic growth of Beltrami type. It is also equivalent to studying the classical Hele-Shaw flow on a Riemann surface, with the Riemannian metric encoding the permeability (see [22]).

There is a vast literature on the Hele Shaw flow (see [20] and the references therein). The main short-time existence and uniqueness result for classical solutions to the Hele-Shaw flow in the case $\kappa \equiv 1$ is due to Kufarev and Vinogradov [40] and dates from 1948. It states that if Ω_0 is a simply connected domain with real analytic boundary, then there exists a unique (indeed real analytic) solution $\{\Omega_t\}_{t\in(-\epsilon,\epsilon)}$ to the Hele-Shaw flow for some $\epsilon > 0$. Observe here that the solution extends both forward and backward in time. In [30] Reissig and von Wolfersdorf gave a new proof of this result using a non-linear version of the Cauchy-Kovalevskaya theorem due to Nishida. Tian [38] provided yet another proof relying on properties of the Cauchy integral of the free boundary, and recently Lin [28] proved the same result using a result of Gustafsson on rational solutions [18] combined with a new perturbation theorem to get to the the general case.

There is also a slightly different setting of the Hele-Shaw flow, where the source of fluid is not a point but a curve inside the starting domain (see e.g. [2, 10, 11]). In [13] Escher and Simonett proved short-time existence of classical solutions to the Hele-Shaw flow in this setting, and in [11, 12] they proved short-time existence of classical solutions to the Hele-Shaw flow with surface tension. There is also related work of Hanzawa [21] on classical solutions to the Stefan problem.

Less has been written about the case of varying permeability. In [22] Hedenmalm-Shimorin proved short-time existence under the assumption that κ is real analytic, and the starting domain has real analytic boundary. They also proved short-time existence when the starting domain is empty, under an additional assumption that $\Delta \log \kappa > 0$, which translates to saying that the Riemann surface in question has negative curvature. See also [23] for related results.

2 Summary of Results

Our first result is the short term existence and uniqueness of the Hele-Shaw flow:

Theorem 2.1. *Let Ω_0 be smooth Jordan domain containing the origin and let κ be a smooth positive function defined in a neighbourhood of $\partial\Omega_0$. Then there exists a smooth positive function κ' defined in a neighbourhood of $\partial\Omega_0$ with $\kappa = \kappa'$ outside of Ω_0, and an $\epsilon > 0$ such that there exists a smooth increasing family of domains $\{\Omega_t\}_{t\in(-\epsilon,\epsilon)}$ which solves the Hele-Shaw flow with varying permeability κ'. The extension κ' is not unique, but a given choice uniquely determines the flow, in the sense that any two such flows agree on their common domain. In particular we have short-time existence and uniqueness of the Hele-Shaw flow with varying permeability forward in time.*

We remark that this goes beyond the results of Hedenmalm-Shimorin since we do not require κ or the domain of the initial domain to be real analytic. In fact if $\kappa \equiv 1$ then it is known the boundary of the domain being real analytic is necessary for there to be a solution backwards in time (see [39] or [19, Thm 10]). Thus part of the above theorem says that this can be circumvented if we modify the permeability appropriately. Moreover if we do make this assumption we can say more and in particular recover Hedenmalm-Shimorin's existence result:

Theorem 2.2. *Suppose in addition to the previous assumptions that κ is real analytic and Ω_0 has real analytic boundary. Then we can take $\kappa' = \kappa$ in a neigh-*

bourhood of $\partial\Omega_0$, and so we have short-time existence and uniqueness of the Hele-Shaw flow both forward and backward in time.

In our opinion, the main interest of this work is not the particulars of the above theorems but the techniques introduced for their proof. We shall show that a solution to the Hele-Shaw flow is equivalent to being able to lift $\Omega_t \subset \mathbb{R}^2 = \mathbb{C}$ to a family of holomorphic discs Σ_t in $\mathbb{C} \times \mathbb{P}^1$ such that

1. Ω_t is the image of the projection of Σ_t to $\mathbb{C}$ and
2. The discs Σ_t attach along their boundaries to a certain totally real submanifold Λ of $\mathbb{C}^2 \subset \mathbb{C} \times \mathbb{P}^1$. and
3. The closure of Σ_t intersects $\mathbb{C} \times \{\infty\} \subset \mathbb{C} \times \mathbb{P}^1$ only at the point $(0, \infty)$.



Figure 3: Lifting the Hele-Shaw flow

This correspondence comes about through thinking of the lifts Σ_t as the graph of certain "Schwarz" functions on Ω_t that are holomorphic except for a simple pole at the origin. Starting with $t = 0$, we construct a smooth strictly subharmonic function ϕ (that encodes the permeability by $\kappa = 1/\Delta\phi$) such that Ω_0 admits such a Schwarz function. Then setting

$$\Lambda = \{(z, \frac{\partial\phi}{\partial z})\} \subset \mathbb{C}^2,$$

the graph of S_0 will be a holomorphic disc Σ_0 whose boundary lies in Λ.

Once this is done we are in a position to apply the well-developed theory of embedded holomorphic discs to prove that the moduli space of embedded discs nearby Σ_0 that are attached to Λ and that pass through $(0, \infty)$ is a smooth manifold of real dimension one which gives both the existence and uniqueness results we require. This dimension count in turn relies on a standard argument using the associated Schottky double and a simple intersection count.

Along the way we use the same technique to analyse the moduli space of quadrature domains (which will be defined in Section 5).

Theorem 2.3. *The moduli space of smooth quadrature domains of order n and connectivity c is a smooth manifold of real dimension $4n + c - 2$.*

It is also interesting to consider the Hele-Shaw flow with empty initial condition. Despite this having a rather natural physical interpretation, there appear to be rather few results in this direction. This may be because many of the techniques in the study of the Hele-Shaw flow rely on identifying the starting domain with the unit disc (say through the Riemann Mapping Theorem) and thus cannot be applied. We shall show how the lifting technique above can be used to prove short term existence in this setting as well, which will require more sophisticated machinery concerning holomorphic discs, specifically results of Donaldson concerning a suitable homogeneous Monge Ampère equation.

Theorem 2.4. *Let κ be a positive smooth function. Then there is an $\epsilon > 0$ and family Ω_t of domains in $\mathbb{C}$ containing the origin that satisfies the Hele-Shaw flow with permeability κ and such that*

$$\mathrm{Area}(\Omega_t) = t \quad \text{for } 0 < t < \epsilon$$

(i.e. the limit of Ω_t as t tends to zero is just the origin).

We remark that this improves on the results of Hedenmalm-Shimorin in that we do not need to assume κ is real analytic or that $\Delta(\log \kappa)$ is positive.

This lifting interpretation of the Hele-Shaw flow through Schwarz functions comes about through looking at the complex moments

$$M_k(t) = \int_{\Omega_t} z^k \frac{1}{\kappa} dA$$

where dA denotes the Lebesgue measure on $\mathbb{C}$. It is known that the Hele-Shaw flow is characterised by the condition that $M_k(t)$ is constant with t for $k \geq 1$ (and so the area $M_0(t)$ is linear with t). One can similarly study the problem of flows Ω_t for which $M_k(t)$ for $k \geq 1$ are allowed to vary in a particular way, and this puts the Hele-Shaw flow into the more general picture of "inverse potential problems". We use the same approach as above to prove a statement about these more general flows:

Theorem 2.5. *Let Ω be a smooth Jordan domain, and V be a smooth, nowhere vanishing outward pointing normal vector field on $\partial\Omega$. Then there exists a smooth variation Ω_t for $t \in [0, \epsilon)$ for some $\epsilon > 0$ such that*

$$\frac{d}{dt}\Big|_{t=0} \partial\Omega_t = V$$

and whose moments vary linearly in t.

Note that this theorem is not immediately obvious since a-priori there could be relations among the higher moments; thus we think of it as showing a kind of independence among them. In fact it is known that if κ is real analytic, and Ω has real analytic boundary, then the moments M_k for $k \geq 0$ provide parameters for the space of domains with analytic boundary near to Ω. We refer the reader to Theorem 7.2 for a more precise version of Theorem 2.5 in this setting. If we set $V = -\nabla p$ in the above theorem, where p denotes the Green function of Ω, we recover the Hele-Shaw flow, and thus this result generalises Theorem 2.1.

Acknowledgements: We first of all wish to thank Björn Gustafsson for conversations and guidance. We also thank Robert Berman, Bo Berndtsson and Claude LeBrun for valuable input, and Johanna Zetterlund for assistance with creation of several images. JR also acknowledges Colin Cotter and Duncan Hewitt for patiently answering basic questions about fluid mechanics. During this project JR was supported by an EPSRC Career Acceleration Fellowship.

Terminology: We identify $\mathbb{R}^2$ with the complex plane $\mathbb{C}$ in the standard way. A *domain* $\Omega \subset \mathbb{R}^2$ is a set that is open and connected and its boundary is $\partial\Omega = \overline{\Omega} - \Omega$ where the bar denotes the topological closure. The *connectivity* of Ω is the number of connected components of $\partial\Omega$. If $\partial\Omega$ can be written locally as the graph of a smooth function then we shall say that Ω has *smooth boundary* or simply that Ω is *smooth*, with analogous definitions if this graph can be taken to be real analytic, and in this case there is a well defined unit outward normal vector field n on $\partial\Omega$. We say Ω is a *smooth Jordan domain* if it is the interior region determined by a smooth Jordan curve (i.e. a smooth non-self intersecting loop in $\mathbb{R}^2$).

If $I \subset \mathbb{R}$ is an interval then a family of domains Ω_t for $t \in I$ is *increasing* if $\Omega_t \subset \Omega_{t'}$ for $t < t'$ and the family Ω_t is said to be *smooth* if the boundary $\partial\Omega_t$ can be written locally as the graph of a smooth function that depends smoothly on t. If Ω_0 is smooth with unit normal vector field n on $\partial\Omega_0$ then an increasing smooth family Ω_t for $t \in (-\epsilon, \epsilon)$ is determined by $\partial\Omega_t = \{x + f(x,t)n_x : x \in \partial\Omega_0\}$ for some positive smooth function $f_t(x) = f(x,t)$ on $\partial\Omega_0$, and the *normal outward velocity* of Ω_t at $t = 0$ is

$$V_0 := \frac{d}{dt}\Big|_{t=0} \partial\Omega_t := \frac{df_t}{dt}\Big|_{t=0} n.$$

3 Lifting the Hele Shaw Flow

3.1 Complex moments

Let Ω_t be a family of domains in $\mathbb{R}^2 = \mathbb{C}$ and κ be a positive smooth function. For $k \geq 0$ the kth *complex moment* of Ω_t is

$$M_k(t) := \int_{\Omega_t} z^k \frac{1}{\kappa} dA,$$

where dA denotes Lebesgue measure. Clearly then $M_0(t)$ is just the area of Ω_t, and we refer to the M_k for $k \geq 1$ as the *higher moments*.

An important discovery by Richardson [31] is that these moments are conserved by the Hele-Shaw flow. Since we are assuming the fluid is injected at a constant rate, we may as well reparametrise t so that $M_0(t) = \mathrm{Area}(\Omega_t) = M_0(0) + t$. The upshot then is that this system is integrable, by which we mean that these moments characterise the Hele-Shaw flow:

Theorem 3.1.

1. *Let Ω_t be given by the Hele-Shaw flow. Then the higher moments $M_k(t)$ remain constant with t.*
2. *Conversely, any smooth increasing family of simply connected domains $\{\Omega_t\}$ for $t \in (-\epsilon, \epsilon)$ that contain the origin and whose higher moments M_k are constant with respect to t is a solution to the Hele-Shaw flow (after a possible reparameterising of t).*

The first statement is Richardson's calculation, namely that if the Hele-Shaw equation (3) is satisfied then

$$\begin{aligned}\frac{d}{dt}M_k(t) &= \frac{d}{dt}\int_{\Omega_t} z^k \frac{1}{\kappa} dA = \int_{\partial\Omega_t} z^k \frac{V_t}{\kappa} ds = -\int_{\partial\Omega_t} z^k \frac{\partial p_t}{\partial n} ds \\ &= \int_{\Omega_t} p_t \Delta(z^k) - z^k(\Delta p_t) dA - \int_{\partial\Omega_t} p_t \frac{\partial z^k}{\partial n}\end{aligned}$$

which vanishes if $k > 0$ and equals 1 when $k = 0$ as $p_t = 0$ on $\partial\Omega_t$ and $\Delta(p_t) = -\delta_0$. The second statement says in effect that this calculation can be reversed (see [20, p22] or [22]).

3.2 Schwarz functions

We next discuss how the moment characterisation of the Hele-Shaw flow can be recast in terms of the existence of Schwarz functions. To describe this, fix a pair (Ω, ϕ) where Ω is a domain in $\mathbb{R}^2 = \mathbb{C}$, and ϕ is a smooth function defined in some neighbourhood of $\partial\Omega$ that is strictly subharmonic (i.e. such that $\Delta\phi$ is positive).

Definition 3.2. We say that a function S is a *Schwarz function* of (Ω, ϕ) if S is a meromorphic function on Ω which extends continuously to $\partial\Omega$ and such that

$$S(z) = \frac{\partial\phi(z)}{\partial z} \quad \text{on } \partial\Omega.$$

Remark 3.3. The model case usually considered is where $\phi = |z|^2$, so $S(z) = \overline{z}$ on $\partial\Omega$. We remark that what we define above is sometimes referred to as an interior Schwarz function, since we are requiring that it be defined on all of Ω. Much of the work of Schwarz functions asks instead that S be defined, and holomorphic, on some neighbourhood of $\partial\Omega$. It turns out, in the model case $\phi(z) = |z|^2$, that such an S exists in this restricted sense if and only if Ω has real analytic boundary (see e.g. the book [37] by Shapiro), and extends to a meromorphic function on Ω if and only if Ω is a quadrature domain [1].

The connection between Schwarz functions in the model case $\phi(z) = |z|^2$ and the Hele-Shaw flow is well known (e.g. [20, 3.7.1]). The following generalises this to suit our needs:

Proposition 3.4. *Let ϕ be a smooth and strictly subharmonic function defined in a domain D. Suppose Ω_t, $t \in (a, b)$, is a smooth increasing family of Jordan domains containing 0 such that for all $t \in (a, b)$, $\partial\Omega_t \subset D$, and for each t there exists a Schwarz function S_t of the pair (Ω_t, ϕ) that is holomorphic on Ω_t except for having a simple pole at 0. Then after some reparametrization $s(t)$, the Ω_s are a solution to the Hele-Shaw flow with varying permeability $\kappa = 1/\Delta\phi$.*

Proof. Pick $t_0 \in (a,b)$ and let $t \in (t_0, b)$. Then for $k \geq 0$ we have,

$$\begin{aligned}\int_{\Omega_t \setminus \Omega_{t_0}} z^k \frac{dA}{\kappa} &= \int_{\Omega_t \setminus \Omega_{t_0}} z^k \Delta\phi dA = \frac{i}{2}\int_{\Omega_t \setminus \Omega_{t_0}} z^k d(\frac{\partial\phi}{\partial z}dz)\\ &= \frac{i}{2}\int_{\partial\Omega_t} z^k \frac{\partial\phi}{\partial z}dz - \frac{i}{2}\int_{\partial\Omega_{t_0}} z^k \frac{\partial\phi}{\partial z}dz\\ &= \frac{i}{2}\int_{\partial\Omega_t} z^k S_t dz - \frac{i}{2}\int_{\partial\Omega_{t_0}} z^k S_{t_0} dz\\ &= \frac{i}{2}\int_{\Omega_t} z^k \bar\partial S_t dz - \frac{i}{2}\int_{\Omega_{t_0}} z^k \bar\partial S_{t_0} dz =\\ &= z^k(0)\pi \mathrm{Res}_0(S_t - S_{t_0}).\end{aligned}$$

Thus the complex moments are preserved, and so Ω_t is a classical solution to the Hele-Shaw flow (3.1) (perhaps after reparametrization). ☐

Thus we wish to prove the existence of Schwarz functions S_t, which is made easier through the freedom of choice in the subharmonic function ϕ. Our next aim is to construct a suitable ϕ that solves this problem for $t = 0$. To guide the subsequent discussion we first recall a classical argument in the case where $\phi = |z|^2$. Consider the Cauchy integral

$$f(z) := \int_{\partial\Omega_0} \frac{\bar\zeta d\zeta}{\zeta - z}.$$

Define f_+ as the restriction of f to Ω_0 and f_- as the restriction of f to the complement of $\overline{\Omega}_0$. Clearly f_+ and f_- are both holomorphic in their respective domain of definition. A careful analysis shows that both f_+ and f_- extend smoothly to $\partial\Omega_0$ and the Sokhotski-Plemelj formula states that on the boundary

$$f_+(z) = \bar z + f_-(z) \tag{4}$$

(see, for example, the book [15] by Gakhov). Moreover it can be shown that the boundary of Ω_0 is real analytic if and only if both f_+ and f_- extend holomorphically to a neighbourhood of $\partial\Omega_0$, and in this case $S := f_+ - f_-$ is holomorphic on this neighbourhood and $S(z) = \overline{z}$ on $\partial\Omega$.

We state first a simple version of what we shall prove:

Proposition 3.5. *Let Ω_0 be a smooth Jordan domain containing 0. Then there exists a smooth strictly subharmonic function ϕ defined on a neighbourhood U of the complement of Ω_0 such that $\Delta\phi = 1$ on $U - \Omega_0$ and so that (Ω_0, ϕ) admits a Schwarz function S_0 that is holomorphic except for having a simple pole at 0.*

For the problem of varying permeability we need something a little stronger:

Proposition 3.6. *Let Ω_0 be a smooth Jordan domain containing 0 and κ be a smooth positive function defined on a neighbourhood of $\partial\Omega_0$. Then there exists a smooth function κ' defined in on a neighbourhood U of $\partial\Omega_0$ such that $\kappa' = \kappa$ on $U - \Omega_0$ and a smooth strictly subharmonic function ϕ defined on U with $\Delta\phi = 1/\kappa'$ such that (Ω_0, ϕ) admits a Schwarz function S_0 that is holomorphic except for having a simple pole at 0.*

Moreover if κ is real analytic, and Ω_0 has real analytic boundary then we can take $\kappa' = \kappa$.

Proof. We need only prove Proposition 3.6. We may as well assume κ is defined on all of $\mathbb{C}$ and is smooth and positive there. Pick some smooth function ϕ_0 on $\mathbb{C}$ such that $\Delta\phi_0 = 1/\kappa$.

Consider the Cauchy integral

$$f(z) := \int_{\partial\Omega_0} \frac{\frac{\partial\phi_0}{\partial z} d\zeta}{\zeta - z}.$$

As above we write f_+ and f_- for the restriction of f to Ω_0 and to the complement of $\overline{\Omega}_0$ respectively. Then the Sokhotski-Plemelj formula says that on $\partial\Omega_0$,

$$f_+(z) = \frac{\partial\phi_0}{\partial z} + f_-(z). \tag{5}$$

Observe next that f_- tends to zero as z tends to infinity. Thus for some constant a we have that

$$f_-(z) = \frac{a}{z} + \tilde{f}(z),$$

where $\tilde{f}$ vanishes to second order at infinity. Such a holomorphic function possesses a $\partial/\partial z$-primitive F on $\mathbb{C} - \overline{\Omega_0}$. Letting $h = 2Re(F)$ we can rewrite equation (5) as

$$f_+(z) - \frac{a}{z} = \frac{\partial}{\partial z}(\phi_0 + h) \quad \text{on } \partial\Omega_0. \tag{6}$$

Now set

$$\phi = \phi_0 + h \quad \text{on } \mathbb{C} - \overline{\Omega}_0,$$

and observe that since h is the real part of a holomorphic function, $\Delta\phi = \Delta\phi_0 = 1/\kappa$. Moreover since f_- had a smooth extension to $\partial\Omega_0$ it follows that ϕ extends smoothly to a neighbourhood of the complement of Ω_0 and we can define $\kappa' = 1/(\Delta\phi)$ which will be positive near $\partial\Omega_0$. Finally $S_0 := f_+(z) - \frac{a}{z}$ is a Schwarz function of (Ω_0, ϕ), which is holomorphic except for a simple pole at 0.

If κ is real analytic and $\partial\Omega_0$ has real analytic boundary, then f_- extends to a holomorphic function over a neighbourhood U of $\partial\Omega_0$ [24, 1.6.4,2.5.1], and h similarly extends to a harmonic function over the boundary. Thus defining ϕ over this neighbourhood as above we have $\Delta\phi = 1/\kappa$ over this neighbourhood as well, and so $\kappa' = \kappa$. ☐

Remark 3.7. From the above proof one sees that in fact the second statement holds under the condition that f_- extends analytically over $\partial\Omega$, the arguments below show that in this case there is a backwards solution to the Hele-Shaw flow.

Remark 3.8. In what follows we will use in an essential way that ϕ is smooth on a neighbourhood of $\partial\Omega_0$. But one should note that the Sokhotski-Plemelj formula holds for much more general domains (e.g. Lipschitz domains) and one still gets a Schwarz function of (Ω_0, ϕ), in this case and ϕ will at least have a $C^{1,\alpha}$ extension.

3.3 Holomorphic Discs

We are now ready to phrase the Hele-Shaw problem in terms of holomorphic discs. Let ϕ be the subharmonic function defined in Proposition 3.6, and S_0 be the Schwarz function for (Ω_0, ϕ). Define

$$\Lambda := \{(z, \frac{\partial\phi}{\partial z})\} \subset \mathbb{C} \times \mathbb{C}.$$

Since this is the graph of a real strictly subharmonic function, it enjoys various properties that will be discussed in the next section (in fact it is a totally real LS-manifold). We shall consider a meromorphic function on an open set in $\mathbb{C}$ as a function to the Riemann sphere $\mathbb{P}^1 = \mathbb{C} \cup \{\infty\}$ in the standard way, giving it the value ∞ at its poles.

Definition 3.9. Suppose (Ω, ϕ) admits a Schwarz function S holomorphic except for having a simple pole at 0. Define

$$\Sigma = \Sigma_\Omega := \{(z, S(z)) \in \mathbb{C} \times \mathbb{P}^1 : z \in \Omega\}. \tag{7}$$

So Σ_Ω is a holomorphic disc in $\mathbb{C} \times \mathbb{P}^1$. Moreover,

1. If π_1 denotes the projection to the first factor then $\pi_1 \colon \Sigma \to \pi_1(\Sigma)$ is an isomorphism,
2. The boundary of Σ lies in Λ,
3. The closure of Σ intersects $\{0\} \times \mathbb{P}^1$ precisely in the point $(0, \infty)$.

Theorem 3.10. *Suppose that Σ_t is a smooth family of holomorphic discs in $\mathbb{C} \times \mathbb{P}^1$ that satisfy the three conditions above and such that $\Omega_t := \pi_1(\Sigma_t)$ is an increasing family of smooth Jordan domains. Then Ω_t is a solution to the Hele-Shaw flow with permeability $\kappa' := 1/(\Delta\phi)$. Moreover any smooth increasing family Ω_t of smooth Jordan domains that solves the Hele-Shaw flow with permeability κ' arises in this way.*

Proof. The first statement just summarises what we have said thus far. Given Σ_t, it can be realised as the graph of a Schwartz function S_t for $\Omega_t := \pi_1(\Sigma_t)$. Condition (3) says that S_t is holomorphic except for having a simple pole at 0. Thus the result we want follows from Proposition 3.4.

For the second statement, suppose that Ω_t is a smooth increasing family of smooth Jordan domains that satisfy the Hele-Shaw flow. Let $p_t(z) = p(z,t)$ be the pressure which is required to satisfies $\Delta p_t = -\delta_0$ on Ω_t and $p_t = 0$ on $\partial\Omega_t$. Now define

$$u_t(z) = u(z,t) = \int_0^t p(z,\tau)d\tau$$

and set

$$S_t = \frac{\partial\phi}{\partial z} - \frac{\partial u_t}{\partial z} - \chi_{\Omega_0}\left(\frac{\partial\phi}{\partial z} - S_0\right),$$

where χ_{Ω_0} denotes the characteristic function of Ω_0. Then $\Delta u = 1 - \chi_{\Omega_0} - t\delta_0$ in the sense of distributions, (see [20, Sec 3.1] or Remark 3.12 below) and one easily checks that S_t is a Schwarz function for Ω_t which is holomorphic except for a simple pole at 0 . Thus the Ω_t are induced from the corresponding family Σ_{Ω_t} as required. ☐

In the following section we show that the moduli space of such holomorphic discs has real dimension 1, and thus our original disc Σ_0 deforms in a smooth one dimensional family Σ_t. Observe that condition (1) and the property that $\pi_1(\Sigma_t)$ is a smooth Jordan domain are open conditions, and thus since they hold for Σ_0 they will hold for Σ_t for small t as well.

Remark 3.11. In much the same way one can interpret the Hele-Shaw flow with multiple inputs, in which case condition (3) above must be suitable modified. We leave the details to the reader.

Remark 3.12. To understand the long-time behaviour of the Hele-Shaw flow, and also for more general starting domains, one has a weak formulation of the problem. Indeed, following Elliott-Janovsky [10], Gustafsson [19] and Sakai [32], [35] it is possible to cast a weak formulation of the Hele-Shaw flow as an obstacle problem.

Let f be a distribution on some domain D. A function u on D is said to solve the corresponding obstacle problem if

$$-\Delta u - f \geq 0, \qquad u \geq 0, \qquad (-\Delta u - f)u = 0. \tag{8}$$

The free boundary $\Gamma(u)$ of the obstacle problem is the boundary of the coincidence $\Lambda(u) := \{x : u(x) = 0\}$. If the domain D has a boundary one also specifies a non-negative function g on ∂D and demand that the solution u should, in addition to satisfying (8), be equal to g on the boundary.

Now let $D = \mathbb{R}^2$ and for any $t > 0$ set

$$f_t := \chi_{\Omega_0} - 1 + t\delta_0.$$

By general potential theory the corresponding obstacle problem has an unique lower semicontinuous solution u_t and one calls the family

$$\Omega_t := \{z : u_t(z) > 0\}$$

the weak solution to the Hele-Shaw flow with initial domain Ω_0. It is know that a smooth solution, if it exists, will coincide with the weak solution, and if the weak solution is regular enough, it is the smooth solution (see e.g. [20]). Moreover there does exists a weak solution that is defined for all time, but in general the Ω_t will eventually be non-smooth or cease to be simply connected.

Now suppose S_0 is a Schwarz function of (Ω_0, ϕ) and suppose that $\Delta\phi = 1$ on the complement of Ω_0. We have seen that such a function exists at least when Ω_0 is Lipschitz. Let Ω_t be the weak solution to the Hele-Shaw flow. We observe that for any $t \in [0, \infty)$, there exists a Schwarz function for (Ω_t, ϕ), holomorphic except for a simple pole at 0. To see this, let u_t be solution to the above obstacle problem described. Then one simply notes that the function

$$S_t := (\frac{\partial \phi}{\partial z} - \frac{\partial u_t}{\partial z}) - \chi_{\Omega_0}(\frac{\partial \phi}{\partial z} - S_0)$$

is the claimed Schwarz function.

Hence using the same lifting we thus see that there is a natural deformation of our initial holomorphic disc Σ_0 that exists for *all* positive time, but whose members need not be smooth and may have more exotic topology. However the possibilities are constrained, as there are strong regularity results due to Sakai [33, 34] for the boundaries $\partial\Omega_t$ of the domains Ω_t belonging to a weak Hele-Shaw flow which says that if $\overline{\Omega}_0 \subseteq \Omega_t$ (which, for example is the case if $\partial\Omega_0$ is C^1) then $\partial\Omega_t$ is real analytic except for having at most a finite number of cusps or double points.

4 Holomorphic curves with boundaries in a totally real submanifold

We have thus seen that the Hele-Shaw flow can be understood from the point of view of Schwarz functions, and in turn in terms of holomorphic discs with boundaries in a given totally real submanifold. There is a rich literature on this topic, we now summarise parts that are relevant to our situation.

4.1 The moduli space

Let M be a complex manifold of complex dimension n. A submanifold Λ is said to be *totally real* if $T\Lambda \cap J(T\Lambda) = \{0\}$, where J denotes the complex structure on M. A totally real submanifold is said to be *maximal* if it has real dimension n.

Let $(\Sigma, \partial\Sigma)$ be a Riemann surface with boundary. A holomorphic non-constant map $f\colon \Sigma \to M$ is called a *holomorphic curve* in M if f extends to a C^1 function on $\partial\Sigma$. If f is an embedding then we call the image of f an *embedded holomorphic curve*. By abuse of terminology we shall refer to this image as Σ, and say that it has boundary in Λ if $f(\partial\Sigma) \subset \Lambda$. In the special case that Σ is the unit disc $D \subset \mathbb{C}$ and $f(\partial D) \subset \Lambda$ we call the image an *embedded holomorphic disc* with boundary in Λ. By the smooth Riemann mapping theorem [7, Thm 3.4], one can equivalently replace the disc D with any smooth Jordan domain.

The moduli space of holomorphic curves with boundary in a totally real maximal submanifold has been studied by numerous authors in different contexts, e.g. Bishop [5], Donaldson [9], Forstnerič [14] and Gromov [16]. In complex analysis this is related to polynomial hulls and the homogeneous Monge-Ampère equation [9], while in symplectic topology it is used in the study of Lagrangian submanifolds [16]. More recently LeBrun and Mason [26, 27] have shown a connection to twistor theory.

The first step in this theory is to give the space of all (not necessarily holomorphic) embedded curves in some regularity class with boundary in Λ the structure of a Banach manifold $\mathcal{B}$. Let Σ be such an embedded curve, let N denote the normal bundle and let E be the normal bundle of $\partial\Sigma$ in Λ. We can identify E with a totally real subbundle of N restricted to $\partial\Sigma$. Locally around Σ the Banach manifold $\mathcal{B}$ is modelled on the Banach space $C^{k+1,\alpha}(\Sigma, N, E)$ of sections of N with boundary values in E (this construction uses the exponential map with respect to some appropriately chosen metric, see [27]).

The next step is to consider a Banach vector bundle $\pi : \mathcal{E} \to \mathcal{B}$ whose fibre over Σ is

$$C^{k,\alpha}(\Sigma, \Lambda^{0,1} \otimes N)$$

and a canonical section $\mathcal{S} : \mathcal{B} \to \mathcal{E}$ whose zero locus $\mathcal{M} := \mathcal{S}^{-1}(0)$ is the space of holomorphic curves. The implicit function theorem for Banach spaces says that $\mathcal{M}$ is a smooth manifold provided that the graph of $\mathcal{S}$ is transverse to the zero section. This is so if and only if at any point Σ in $\mathcal{M}$, the linearisation D_Σ, which by definition is $D\mathcal{S}$ composed with the projection to the fibre of $\mathcal{E}$, is surjective and has a right inverse. Now D_Σ is nothing but the Cauchy-Riemann operator $\bar\partial$ that maps sections of N with boundary values in E to $(0,1)$-forms with values in N. It is well-known that this operator is Fredholm, so what remains to establish transversality of $\mathcal{S}$ is surjectivity. Thus if the Cauchy-Riemann operator is surjective, then $\mathcal{M}$ is a smooth manifold and its dimension is given by the index of $\bar{\partial}$.

In general the problem of proving transversality involves picking a new generic almost complex structure on the manifold M. However for our purposes M has complex dimension 2 and then the problem becomes significantly easier and, as we shall see, reduces to the computation of a single topological invariant of a related holomorphic curve.

4.2 Dimension count and the Schottky double

Following LeBrun in [27], we now discuss how to use the Schottky double to compute the dimension of the moduli space $\mathcal{M}$ of embedded holomorphic curves in a manifold M of complex dimension 2.

Let Σ be a compact Riemann surface with boundary $\partial\Sigma$. By gluing it together with its complex conjugate $\bar{\Sigma}$ along $\partial\Sigma$ we get a closed Riemann surface $\mathcal{X}$ (assuming some regularity of Σ, see [27]). In the theory of quadrature domains $\bar{\Sigma}$ is called the Schottky double of Σ [17]. The complex conjugate $\bar{N}$ of N is a holomorphic vector bundle on $\bar{\Sigma}$, and by gluing N and $\bar{N}$ along $\partial\Sigma$ so that $E = \bar{E}$ we get a holomorphic vector bundle $\mathcal{N}$ on $\mathcal{X}$.



Figure 4: The Schottky double

There is an antiholomorphic involution ρ on $\mathcal{X}$ which lifts to $\mathcal{N}$ such that E is the fixed point set of ρ. As noted in [27], this implies that E is real analytic as a subspace of $\mathcal{N}$. This involution thus acts on each of the vector spaces

$$C^{k+1,\alpha}(\Lambda, \mathcal{O}(\mathcal{N})) \text{ and } H^p(\mathcal{X}, \mathcal{O}(\mathcal{N})) \text{ for } p = 0, 1.$$

Let $C^{k+1,\alpha}_\rho(\mathcal{X}, \mathcal{O}(\mathcal{N}))$ and $H^p_\rho(\mathcal{X}, \mathcal{O}(\mathcal{N}))$ denote the invariant subspaces. Thus

$$H^p(\mathcal{X}, \mathcal{O}(\mathcal{N})) = H^p_\rho(\mathcal{X}, \mathcal{O}(\mathcal{N})) \oplus iH^p_\rho(\mathcal{X}, \mathcal{O}(\mathcal{N})).$$

LeBrun shows in [27] that

$$C^{k+1,\alpha}(\Sigma, N, E) \cong C^{k+1,\alpha}_\rho(\mathcal{X}, \mathcal{O}(\mathcal{N}))$$

with a canonical isomorphism. Also [27, Lemma 1] the kernel of the Cauchy-Riemann operator is canonically isomorphic to $H^0_\rho(\mathcal{X}, \mathcal{O}(\mathcal{N}))$ while the cokernel is canonically isomorphic to $H^1_\rho(\mathcal{X},\mathcal{O}(\mathcal{N}))$. Thus $\bar\partial$ is surjective if and only if $h^1(\mathcal{X},\mathcal{O}(\mathcal{N}))=0$, and then the index is equal to $h^0(\mathcal{X},\mathcal{O}(\mathcal{N}))$.

Now suppose that when the complex dimension of the ambient manifold M is two. Then $\mathcal{N}$ is a complex line bundle, and so as long as $\deg\mathcal{N}>2g-2$, where g is the genus of $\mathcal{X}$, then the numbers $h^0(\mathcal{X},\mathcal{O}(\mathcal{N}))$ and $h^1(\mathcal{X},\mathcal{O}(\mathcal{N}))$ are topological invariants that only depend on g and the degree of $\mathcal{N}$.

If Λ happens to be the fixed point set of an antiholomorphic involution ρ of M, then the Schottky double $\mathcal{X}$ of Σ can be identified with the Riemann surface $\Sigma\cup\rho(\Sigma)$, and $\mathcal{N}$ is then identified with the normal bundle of $\Sigma\cup\rho(\Sigma)$ in M. Hence a strategy to compute the numbers $h^0(\mathcal{X},\mathcal{O}(\mathcal{N}))$ and $h^1(\mathcal{X},\mathcal{O}(\mathcal{N}))$ for a general pair (Σ,Λ) in the four dimensional case is to continuously perturb (Σ,Λ) to some (Σ',Λ') where Λ' is the fixed point set of an involution ρ, and then use the adjunction formulae to compute the degree of the normal bundle of $\Sigma'\cup\rho(\Sigma')$.

5 Moduli spaces of quadrature domains

We digress in this section from the Hele-Shaw flow, and consider the above setup in the case $\Lambda=\{(z,\bar z)\}$ which is related the concept of quadrature domains.

Definition 5.1. A bounded domain Ω in $\mathbb{C}$ is called a *quadrature domain* if there exist finitely many points $a_1,...,a_m\in\Omega$ and coefficients c_{kj} such that for any integrable holomorphic function f on Ω the quadrature identity holds:

$$\int_\Omega f dA=\sum_{k=1}^m\sum_{j=0}^{n_k-1}c_{kj}f^{(j)}(a_k).$$

The integer $n=\sum_{k=1}^m n_k$ is called the *order* of the quadrature domain (assuming $c_{k,n_k-1}\neq 0$). We say that a quadrature domain is smooth if its boundary is smooth. We also recall that the *connectivity* c of a domain Ω is the number of connected components of $\partial\Omega$.

Assume that $(\Omega,\phi(z)=|z|^2)$ has a Schwarz function S. Then

$$\begin{aligned}\int_\Omega f dA=\frac{i}{2}\int_\Omega fd(\bar z dz) &= \frac{i}{2}\int_{\partial\Omega} f\bar z dz\\ &= \frac{i}{2}\int_{\partial\Omega} fSdz=\frac{i}{2}\int_\Omega f\bar\partial S dz\\ &= \sum_{k=1}^m\sum_{j=0}^{n_k-1}c_{kj}f^{(j)}(a_k),\end{aligned}$$

where a_k are the poles of S and n_k is the order of the pole at a_k, and thus Ω is a quadrature domain. It is proved in [1, Lemma 2.3] that the converse holds as well. Clearly then the order of the domain n equals the number of poles of S counted with multiplicity.

It is also clear that Schwarz function S is uniquely determined by the domain, since if we had two Schwarz functions, their difference would be a meromorphic function that was zero on the boundary, and therefore this difference must be zero.

From now on assume that the quadrature domain Ω has smooth boundary. Consider the graph

$$\Sigma:=\{(z,S(z)):z\in\Omega\}$$

as an embedded holomorphic curve in $\mathbb{C} \times \mathbb{P}^1$. It is attached along its boundary to the totally real submanifold

$$\Lambda = \{(z, \bar{z}) : z \in \mathbb{C}\}.$$

Since Λ is the fixed point set of the antiholomorphic involution $\rho : (z, w) \mapsto (\bar{w}, \bar{z})$, the Schottky double $\mathcal{X}$ is naturally identified with $\Sigma \cup \rho(\Sigma)$ and the normal bundle $\mathcal{N}$ is identified with the normal bundle of $\Sigma \cup \rho(\Sigma)$.

We wish to calculate the degree of $\mathcal{N}$. By the adjunction formula,

$$\mathcal{N} = \mathcal{O}(\mathcal{X})_{|\mathcal{X}} = (K_{\mathbb{P}^1 \times \mathbb{P}^1})^*_{|\mathcal{X}} \otimes K_{\mathcal{X}}$$

where $\mathcal{O}(\mathcal{X})$ denotes the line bundle associated to the divisor $\mathcal{X}$. The line bundle $K_{\mathbb{P}^1 \times \mathbb{P}^1}$ is represented by the divisor $-2(H_1 + H_2)$ where H_1 and H_2 denote the hyperplanes $\mathbb{P}^1 \times \{\infty\}$ and $\{\infty\} \times \mathbb{P}^1$ respectively. Thus

$$\deg \mathcal{N} = 2(H_1 + H_2) \cdot \mathcal{X} + \deg K_{\mathcal{X}}.$$

where the dot represents the intersection number of the relevant divisors. Clearly

$$H_1 \cdot \mathcal{X} = H_2 \cdot \mathcal{X} = n,$$

and

$$\deg K_{\mathcal{X}} = 2g - 2,$$

where g denotes the genus of $\mathcal{X}$. Combined this yields that

$$\deg \mathcal{N} = 4n + 2g - 2.$$

Now $n > 0$, so $\deg \mathcal{N} \geq 2g + 2$ which implies $h^1(\mathcal{X}, \mathcal{N}) = 0$, and by Riemann-Roch that

$$h^0(\mathcal{X}, \mathcal{N}) = \deg \mathcal{N} - g + 1 = 4n + g - 1.$$

Finally note that the genus g is precisely $c - 1$, where c is the connectivity number of Ω.

Hence the moduli space of embedded holomorphic curves with boundaries in Λ is a smooth manifold of real dimension $4n + c - 2$ near Σ. Any such holomorphic curve lying close enough to Σ gives a Schwarz function of a smoothly bounded quadrature domain. By the fact that all quadrature domains have unique Schwarz functions (again with respect to $\phi(z) = |z|^2$), it follows that the moduli space of smoothly bounded quadrature domains of order $n > 0$ and connectivity g is a manifold of real dimension $4n + c - 2$. We have thus proved Theorem 2.3 as stated in the introduction.

6 Short-time existence of the Hele-Shaw flow

This section completes the proof of Theorem 2.1. To ease notation let $\Omega = \Omega_0$ be a smooth Jordan domain. We saw above that we could construct a generalised Schwarz function S of the pair (Ω, ϕ) where $\Delta\phi = 1/\kappa$ on the complement of Ω and ϕ extends smoothly to a neighbourhood of the complement. If Σ denotes the graph of S and Λ the graph of $\frac{\partial \phi}{\partial z}$ we have that Σ is a holomorphic disc whose boundary lies in the totally real submanifold Λ.

We want to analyse the moduli space of holomorphic discs near Σ with boundary in Λ that intersects the hyperplane $H_1 = \mathbb{P}^1 \times \{\infty\}$ at the point $(0, \infty)$.

First we consider the moduli space of all such discs, irrespective of their intersection with H_1. We need to calculate the numbers $h^0(\mathcal{X}, \mathcal{N})$ and $h^1(\mathcal{X}, \mathcal{N})$. As we continuously deform the domain Ω to the unit disc Ω', the pair $(\mathcal{X}, \mathcal{N})$ and submanifold Λ continuously deform to the pair $(\mathcal{X}', \mathcal{N}')$ and to $\Lambda' = \{(z, \bar{z}) : z \in \mathbb{C}\}$ respectively. Thus the Schwarz function for Ω' is simply $\frac{1}{z}$, i.e. the usual (interior) Schwarz function of the unit disc thought of as a quadrature domain. Since the numbers $h^0(\mathcal{X}, \mathcal{N})$ and $h^1(\mathcal{X}, \mathcal{N})$ are invariant under continuous deformation, the calculation in the previous section yields that $h^1(\mathcal{X}, \mathcal{N}) = 0$ and $h^0(\mathcal{X}, \mathcal{N}) = 3$, since $n = 1$ and $g = 0$. Thus the required moduli space is a smooth manifold of real dimension 3 near Σ.

Considering moduli spaces of holomorphic curves passing through a given point is also standard in the theory (see, for example, [29]). We simply replace our Banach manifold $\mathcal{B}$ with the subspace of curves going through the point, which is locally modelled on the Banach subspace of sections to the normal bundle that vanish at that point. We will denote this space by $C^{k+1,\alpha}(\Sigma, N, E)_0$. Following [27] it follows that this space is naturally isomorphic to the space of ρ invariant sections of $\mathcal{N}$ that vanish at both $(0, \infty)$ and $\rho((0, \infty))$. We denote this space by $C^{k+1,\alpha}_\rho(\mathcal{X}, \mathcal{N})_0$. The section $\mathcal{S}$ and its linearisation $\bar{\partial}$ are simply the restrictions of the original ones.

We check first that $\bar{\partial}$ is still surjective. Pick an arbitrary element

$$g \in C^{k,\alpha}(\Sigma, \Lambda^{0,1} \otimes N).$$

By the above there exists a section $f \in C^{k+1,\alpha}(\Sigma, N, E)$ such that $\bar{\partial} f = g$. Since the degree of $\mathcal{N}$ is positive and $\mathcal{X}$ is isomorphic to $\mathbb{P}^1$ we know that there is a holomorphic section, s say, of $\mathcal{N}$ that is equal to f at $(0, \infty)$ but zero at $\rho((0, \infty))$. Then the section $s + \rho(s)$ lies in $H^0_\rho(\mathcal{X}, N)$, and thus corresponds to an element $\tilde{s}$ in the kernel of $\bar{\partial}$. We have that $\tilde{s} = f$ at the point $(0, \infty)$. Thus $f - \tilde{s}$ lies in $C^{k+1,\alpha}(\Sigma, N, E)_0$ and $\bar{\partial}(f - \tilde{s}) = g$. This shows that the restriction of $\bar{\partial}$ is still surjective.

The kernel of the restricted Cauchy-Riemann operator is of course the intersection of the original kernel with the subspace $C^{k+1,\alpha}(\Sigma, N, E)_0$. Under the natural isomorphism this equals the intersection of $H^0_\rho(\mathcal{X}, N)$ and $C^{k+1,\alpha}_\rho(\mathcal{X}, \mathcal{N})_0$. Let $H^0(\mathcal{X}, N)_0$ denote the subspace of holomorphic sections that vanish at both $(0, \infty)$ and $\rho((0, \infty))$. The involution ρ acts on this space by

$$\rho(s)(x) := \rho(s(\bar{\rho}(x))),$$

and if $H^0_\rho(\mathcal{X}, N)_0$ denotes the invariant real subspace we have that

$$H^0(\mathcal{X}, N)_0 \equiv H^0_\rho(\mathcal{X}, N)_0 \oplus i H^0_\rho(\mathcal{X}, N)_0. \tag{9}$$

Clearly $H^0(\mathcal{X}, N)_0$ is the intersection of $H^0_\rho(\mathcal{X}, N)$ and $C^{k+1,\alpha}_\rho(\mathcal{X}, \mathcal{N})_0$, and from (9) we get that the real dimension of $H^0_\rho(\mathcal{X}, N)_0$ is equal to the complex dimension of $H^0(\mathcal{X}, N)_0$. Since the degree of $\mathcal{N}$ is two and $\mathcal{X} \cong \mathbb{P}^1$ the complex dimension of $H^0(\mathcal{X}, N)_0$ is one. Thus the index of the restricted Cauchy-Riemann operator is one.

Together with the surjectivity this implies that the moduli space of nearby holomorphic discs with boundary in Λ and going through $(0, \infty)$ is a smooth manifold of real dimension one. Thus $\Sigma_0 = \Sigma$ varies in a smooth family Σ_t for t sufficiently small, and by Theorem 3.10 it remains to prove that the family of domains $\Omega_t = \pi(\Sigma_t)$ is increasing.

The tangent space of the family of discs at Σ is given by $H^0_\rho(\mathcal{X}, N)_0$ restricted to Σ. Since a non-zero element in $H^0_\rho(\mathcal{X}, N)_0$ vanishes only at the points $(0,\infty)$ and $\rho((0,\infty))$ it is nonvanishing along the boundary of Σ. The real subspace E is one-dimensional and is has one direction going out of Ω_0 and one direction going in. The nonvanishing implies that a nonzero tangent points either outwards everywhere or inwards everywhere. Thus (replacing t with $-t$ if necessary) the family Ω_t is increasing.

7 The inverse potential problem and moment flows

Let ρ be a density function on $\mathbb{C}$ (so we have changed notation from previous sections in which this density was played by $1/\kappa$). The associated potential of a domain Ω is the subharmonic function

$$U^\Omega(z) := \int_\Omega \log|z-w|^2 \rho(w) dA(w)$$

where dA is the Lebesgue measure. Note that U^Ω is harmonic on the complement of Ω. Let D be some big disc containing Ω. The inverse potential problem asks whether a given harmonic function U on the complement of D (with proper growth at infinity) can be realised (uniquely) as the potential of a domain Ω in D.

To see the relationship with the complex moments consider

$$G := \frac{\partial U}{\partial z},$$

which is holomorphic if U is harmonic. If $U = U^\Omega$ for some domain Ω then

$$G(z) = \frac{\partial}{\partial z}\int_\Omega \log|z-w|^2 \rho(w) dA(w) = \int_\Omega \frac{\rho(w)}{z-w} dA(w). \tag{10}$$

Now for large $|z|$ we can write this integral as a convergent power series in z^{-k},

$$\int_\Omega \frac{\rho(w)}{z-w} dA(w) = \sum_{k=1}^\infty a_k z^{-k},$$

where

$$a_k := \int_\Omega w^{k-1}\rho(w) dA(w).$$

Thus the coefficients a_k of the power series of G at infinity equals the complex moments of Ω with respect to the measure ρdA. Thus the inverse potential problem be restated as asking for a (unique) domain with prescribed complex moments.

This potential problem has a rich history, both for the plane and for higher dimensions, and also for more complicated kernels replacing the expression $\ln|z-w|^2$ above (see, for example, [24] and the references therein). For this particular problem in the plane it is known that prescribing the complex moments does not uniquely determine the domain (although it appears to be unknown if this uniqueness can fail for domains with smooth boundary). However starting with a suitable smooth domain Ω_0, for small variations U_t of the potential U^{Ω_0} there is a unique Ω_t close to Ω such that $U_t = U^{\Omega_t}$. Thus one can think of the set of moments as giving local parameters for the space of domains. We refer the reader to [24, Sec 2.4] and [6, Sec. 2] for precise statements. We also point out the interesting work of Wiegmann and Zabrodin in this context, which connects these parameters to an integrable system coming from the dispersionless Toda lattice [41].

We have seen that the Hele-Shaw flow can be phrased as a local existence question related to the inverse potential problem. Namely, if Ω is a smooth Jordan domain containing the origin, then there is a non-trivial one parameter family of smooth domains Ω_t, all having the same higher moments as Ω and whose area increases linearly. In this section we will prove short-time existence of a similar flow in which all the higher moments are allowed to vary linearly. Variants of the Hele-Shaw flow under some other force (e.g. gravity) are considered by Escher-Simonett [11, 12, 13] and this gives a physical interpretation to some of these variations.

Theorem 7.1. *Let Ω be a smooth Jordan domain, and V be a smooth, nowhere vanishing outward pointing normal vector field on $\partial\Omega$. Then there exists a smooth variation Ω_t for $t \in [0,\epsilon)$ for some $\epsilon > 0$ such that*

$$\frac{d}{dt}\Big|_{t=0}\partial\Omega_t = V$$

whose moments vary linearly in t.

Note that without the assumption on the vector field V pointing outwards this theorem is not true, for example in the Hele-Shaw flow when $\rho \equiv 1$ and $V = \nabla p$, then this would mean solving the Hele-Shaw flow backwards in time, which is impossible unless the boundary of Ω is real analytic.

Of course the way in which the moments vary will depend on V. To discuss this suppose V is any normal vector field defined on the boundary of a smooth Jordan domain Ω. Write the measure $V\rho ds$ as the restriction of a complex one form $g(z)dz$ to the curve $\partial\Omega$. The complex-valued function g is then uniquely determined on $\partial\Omega$. Consider the Cauchy integral

$$f = f^{\Omega,V} = \int_{\partial\Omega} \frac{g(w)}{z-w}dw,$$

and as above let $f_+ = f_+^{\Omega,V}$ denote the restriction of f to Ω and $f_- = f_-^{\Omega,V}$ the restriction to the complement of $\overline{\Omega}$. Recall that both f_+ and f_- has a smooth extension to the boundary and

$$f_+ = g + f_- \quad \text{on } \partial\Omega.$$

Now if Ω_t is a smooth variation of Ω with initial direction V, the variation of the complex moments $M_k(t)$ of Ω_t satisfy

$$\frac{dM_k}{dt}\Big|_{t=0} = \int_{\partial\Omega} z^k V\rho ds.$$

Since f_+ is holomorphic in Ω it then follows that

$$\frac{dM_k}{dt}|_{t=0} = -\int_{\partial\Omega} z^k f_-(z)dz.$$

Thus writing f_- as a power series in a neighbourhood of infinity,

$$f_- = \sum_{k=1}^{\infty} b_k z^{-k}$$

we conclude

$$\frac{dM_k}{dt}|_{t=0} = -2\pi i b_{k+1}.$$

We think of the condition that f_- extend to an analytic function across $\partial\Omega$ as a kind of "growth condition" on the coefficients b_k that determine the variation of the moments. With this said we can prove something more precise:

Theorem 7.2. *Let Ω be a smooth Jordan domain. Suppose complex coefficients b_k for $k \geq 0$ are given with $ib_0 \in \mathbb{R}$, so that the function $f_- = \sum_k b_k z^{-k}$ defined in a neighbourhood of infinity extends to a holomorphic function across $\partial\Omega$. Then there exists a smooth density function ρ' with $\rho' = \rho$ outside of Ω and a smooth family $\{\Omega_t\}_{t\in(-\epsilon,\epsilon)}$, whose moments $M_k(t)$ taken with respect to the measure $\rho' dA$ vary linearly as*

$$M_k(t) = M_k(0) - 2\pi i b_{k+1} t.$$

If moreover Ω has real analytic boundary and ρ is real analytic then we can take $\rho' = \rho$.

Remark 7.3. This theorem gives one precise way to state that these moments form parameters for the space of domains near Ω. The final statement, under the extra assumption that the boundary is real analytic, appears to be well known and to have been proved in a number of contexts; it follows, for instance, from [6, Sec. 2]. We are not aware of a similar statement in the smooth case, or one such as Theorem 7.1 that uses the extra data that the initial normal vector field is outward pointing.

The proof starts with the following uniqueness result:

Proposition 7.4. *Let V_1 and V_2 be two smooth normal vector fields defined on the boundary of Ω. If for all $k \geq 0$ we have that*

$$\int_{\partial\Omega} z^k V_1 \rho ds = \int_{\partial\Omega} z^k V_2 \rho ds, \tag{11}$$

then it follows that $V_1 = V_2$.

Proof. Let g_1 and g_2 be the complex valued functions so that $V_i \rho ds = g_i dz$ on $\partial\Omega$ for $i = 1, 2$. By our calculations above it follows from (11) that $g_1 - g_2$ extends holomorphically to Ω. Let u be a conformal map from the unit disc to Ω and consider the pullback of $(g_1 - g_2)dz$ to the unit disc. It will be a holomorphic one form which can be written as $h(w)dw$, where w denotes the holomorphic coordinate on the unit disc. The fact that $(g_1 - g_2)dz$ restricts to a real measure on $\partial\Omega$ implies that the restriction of $h(w)dw$ to the unit circle is real. Thus in turn implies that $wh(w)$ is real valued on the unit circle. It follows that $wh(w) \equiv 0$ and thus $g_1 = g_2$, and consequently $V_1 = V_2$. □

Recall that we have a totally real submanifold Λ which is the graph of $\frac{\partial \phi}{\partial z}$ where $\Delta \phi = \rho$ outside of Ω. We also have a holomorphic disc Σ attached to Λ which projects down to Ω. As we saw in Section 6, Σ belongs to a one parameter family of holomorphic discs attached to Λ which goes through the point $(0, \infty)$, and the projection of these yields the Hele-Shaw flow (the permeability being $1/\rho$).

We record the following theorem.

Theorem 7.5. *Let Σ and Λ be as above. Given any small smooth perturbation Λ_s of Λ, the family Σ_t deforms to a smooth family $\Sigma_{s,t}$ of holomorphic discs going through $(0, \infty)$ attached to Λ_s along their boundaries.*

Proof. This is standard in the theory of moduli spaces of embedded holomorphic curves (see e.g. [27]). Let Λ_s be a smooth parametrised perturbation of Λ, $s \in (-\epsilon, \epsilon)$. The Banach space $\mathcal{B}$ one considers is the space of embedded discs in some regularity class together with the parameter s, where the discs passes through $(0, \infty)$ and is attached to Λ_s. The Banach bundle $\mathcal{E}$ is defined just as before, and so is the section $\mathcal{S}$. The linearisation of S is the $\bar{\partial}$ operator which maps the tangent space of $\mathcal{B}$ to the fibre of $\mathcal{E}$. Let Σ_s be a smooth family of smooth discs lying in $\mathcal{B}$ such that $\Sigma_0 = \Sigma$ and Σ_s attaches to Λ_s. Then the tangent space of $\mathcal{B}$ is spanned by $C^{k+1,\alpha}(\Sigma, N, E)_0$ together with the derivative of Σ_s at $s = 0$, which we will denote by f. We have already established that the linearisation $\bar{\partial}$ is surjective, even when restricted to the subspace $C^{k+1,\alpha}(\Sigma, N, E)_0$. Therefore there is a section g in $C^{k+1,\alpha}(\Sigma, N, E)_0$ such that $\bar{\partial} f = \bar{\partial} g$. Thus the kernel of $\bar{\partial}$ is spanned by the kernel of the restricted $\bar{\partial}$ operator together with $f - g$, so it has dimension 2. By the implicit function theorem for Banach manifolds the space of holomorphic discs nearby Σ lying in $\mathcal{B}$ is a manifold of real dimension 2. Also the subsets which are attached to Λ_s for a given s will be one dimensional submanifolds. □

We now prove Theorem 7.1. Let f_- be the restriction to the complement of $\bar{\Omega}$ of

$$f = \int_{\partial \Omega} \frac{g(w)}{z - w} dw,$$

where $V \rho ds = g(z) dz$, and choose some smooth extension to a neighbourhood U of the boundary of Ω. We will consider the smooth perturbation of Λ given by

$$\Lambda_s := \{(z, \frac{\partial \phi}{\partial z} + s f_-) : z \in U \supset \partial \Omega\}.$$

By Theorem 7.5 the family of holomorphic discs corresponding to the Hele-Shaw flow Σ_t perturbs to a smooth family $\Sigma_{s,t}$ with boundary in Λ_s. For each s select the holomorphic disc $\Sigma'_s := \Sigma_{s,t(s)}$ so that the corresponding Schwartz function S_s has the same residue as S_0. That Σ'_s is a smooth family follows from the ordinary implicit function theorem in finite dimensions. Let Ω_s be the projection of Σ'_s, we have for small s,

$$\int_{\Omega_s} z^k \rho dA - \int_{\Omega_0} z^k \rho dA = \int_{\partial \Omega_s} z^k \frac{\partial \phi}{\partial z} dz - \int_{\partial \Omega_0} z^k \frac{\partial \phi}{\partial z} dz =$$

$$= \int_{\partial \Omega_s} z^k (S_s - s f_-) dz - \int_{\partial \Omega_0} z^k S_0 dz = -s \int_{\partial \Omega_s} z^k f_- dz. \tag{12}$$

Here we used the fact that $S_s = \frac{\partial \phi}{\partial z} + s f_-$ on $\partial \Omega_s$ and that S_s and S_0 have the same residue at 0.

If V_0 is the initial normal vector field on $\partial\Omega$ corresponding to the perturbation Ω_s, then (12) implies that V_0 and V induce the same initial variation of moments. From Proposition 7.4 it thus follows that the initial direction of Ω_s is given by V. In particular this initial direction is, by assumption, non-vanishing and outward pointing, so by smoothness $\partial\Omega_s$ will lie in the complement of $\bar{\Omega}$ for small s. But on this complement f_- is holomorphic, and

$$\int_{\partial\Omega_s} z^k f_- dz = 2\pi i b_{k+1}$$

(which in particular is independent of s). Thus by (12) again, all the moments of Ω_s vary linearly as required.

The proof of Theorem 7.2 is identical, only now since we do not have the hypothesis that the flow is outward initially we must take the moments with respect to $\rho' = \Delta\phi$ which is only equal to ρ outside of Ω unless real analytic is assumed (as in Proposition 3.6)

8 The Hele-Shaw flow and the complex HMAE

In this section we will describe a connection between the Hele-Shaw flow and a certain complex Homogeneous Monge-Ampère Equation (HMAE). We will use this connection in order to prove short-time existence of the Hele-Shaw flow with varying permeability with empty initial condition. We recall that this extends a theorem of Hedenmalm-Shimorin [22] who prove this under the assumption κ is real analytic and $\log\kappa$ is strictly subharmonic.

8.1 The Homogeneous Monge-Ampère Equation

The complex Homogeneous Monge-Ampère Equation is a higher dimensional analogue of the Laplace equation on complex manifolds. We first describe some work of Donaldson [9] that gives an equivalence between regular solutions of a certain HMAE and families of holomorphic discs with boundaries in a so called "LS submanifold" which will be defined below.

Let X be a compact Kähler manifold of complex dimension n, and let ω_0 be a Kähler form. This means that locally ω_0 is equal to $i\partial\bar{\partial}\phi$ where ϕ is a smooth strictly plurisubharmonic function. For example, if X has an ample line bundle L, and h is a positive metric on L, then the curvature form $i\Theta_h$ is a Kähler form on X which lies in the first Chern class of L. Any Kähler class in the same cohomology class as ω_0 can be written as $\omega_0 + i\partial\bar{\partial}\phi$ where ϕ is a global smooth function.

Let D denote the closed unit disc and $D^\times = D - \{0\}$. Also let Ω_0 be the pullback of ω_0 by the projection $\pi\colon X \times D \to D$. Suppose next that F is a smooth function on $X \times \partial D$. We shall write $F_\tau(\cdot) = F(\cdot, \tau)$ thought of as a function on X, and suppose that $\omega_0 + i\partial\bar{\partial}F_\tau$ is a Kähler form for all $\tau \in \partial D$. A regular solution to the associated HMAE is an extension of F to a smooth function Φ on $X \times D$ such that $\omega_0 + i\partial\bar{\partial}\Phi_\tau$ is a Kähler form for all $\tau \in D$ and

$$(\Omega_0 + i\partial\bar{\partial}\Phi)^{n+1} = 0$$

on $X \times D$.

Now suppose Φ is a regular solution to the above HMAE. Then kernel of $\Omega_0 + i\partial\bar{\partial}\Phi$ defines a one dimensional complex distribution, which one can show is integrable, thus giving a foliation of $X \times D$ by holomorphic discs transverse to the X fibres (see [3] and [9]).

A further connection with holomorphic discs comes about through the following idea due to Donaldson [9]. Let $\Theta := \Theta_1 + i\Theta_2$ denote a holomorphic 2-form on a complex manifold W, where Θ_1 and Θ_2 are real symplectic forms. A (real) submanifold Y of W is called a *LS submanifold* if it is Lagrangian with respect to Θ_1 while it is symplectic with respect to Θ_2.

We next sketch the construction of a holomorphic fibre bundle W_X over X such that Kähler forms in the same class as ω_0 are described as closed LS submanifolds in W_X. If ω_0 has a potential ϕ_U in some open set U, we identify W_U with the $(1, 0)$ part of the complex cotangent bundle. If z_i are local holomorphic coordinates any $(1, 0)$-form can be written as $\sum_i \xi_i dz_i$, thus (z_i, ξ_j) are local holomorphic coordinates of W_U. We also have non-degenerate holomorphic 2-form, namely $\Theta := \sum_i d\xi_i \wedge dz_i$.

If V is some other open set where ω_0 has a potential ϕ_V, then on $U \cap V$ the transition function is given by $\partial(\phi_V - \phi_U)$. It means that the section, locally defined as $\partial\phi_U$, is in fact globally defined. By a simple calculation the graph of this section turns out to be a LS submanifold in W_X. Let ω be some other Kähler form in the same class as ω_0. Then $\omega = \omega_0 + i\partial\bar{\partial}\psi$ for some smooth global function ψ, and the section $\partial\phi_U + \partial\psi$ defines a section of W_X. By the same reasons this is also a LS submanifold. Donaldson shows that Kähler forms in the class of ω_0 are in one-to-one correspondence with the closed LS submanifolds of W_X.

Now let F be a smooth function on $X \times \partial D$ such that $\omega_0 + i\partial\bar{\partial}F_\theta$ is positive for $\theta \in \partial D$. For each $\theta \in \partial D$, denote by Λ_θ denote the corresponding LS submanifold in W_X.

Theorem 8.1 (Donaldson [9]). *There is a regular solution to the complex HMAE with boundary values given by F iff there is a smooth family of holomorphic discs $g_x : D \to W_X$ parametrised by $x \in X$ with*

1. $\pi(g_x(0)) = x \in X$;
2. *for each $\theta \in \partial D$ and each $x \in X$,*
$$g_x(\theta) \in \Lambda_\theta;$$
3. *for each $\tau \in D$ the map $x \mapsto \pi(g_x(\tau))$ is a diffeomorphism from X to X.*

In fact for fixed $\tau \in D$, the image of the map $x \mapsto g_x(\tau)$ is the LS submanifold Λ_τ associated to the Kähler form $\omega_0 + i\partial\bar{\partial}\Phi_\tau$.

Using this theorem Donaldson applies the theory of the moduli spaces for embedded holomorphic discs with boundaries in a totally real submanifold to deduce the following:

Theorem 8.2 (Donaldson [9]). *The set of boundary conditions F for which the HMAE above has a regular solution Φ is open in the C^2 topology.*

8.2 Short-time existence of the Hele-Shaw flow with empty initial condition

We will now consider a HMAE as above with $X = \mathbb{P}^1$ and $\omega_0 = \omega_{FS}$, the Fubini-Study form on $\mathbb{P}^1$. If we in the usual way think of $\mathbb{P}^1$ as $\mathbb{C} \cup \{\infty\}$, then on $\mathbb{C}$ the

Fubini-Study form has local potential $\phi_{FS} = \log(1+|z|^2)$. The local potential of a Kähler form in the same class as ω_{FS} is a smooth strictly subharmonic ϕ such that $\phi - \log(1+|z|^2)$ has a smooth extension to the point at infinity.

Now fix such a potential ϕ with $\Delta\phi = 1/\kappa$, where κ is a positive smooth function that encodes the permeability for the Hele-Shaw flow. For any $\theta \in \partial D$ set

$$F(z,\theta) := \phi(\theta z) - \phi_{FS}(z). \tag{13}$$

Under the natural identification of $W_{\mathbb{C}}$ with $\mathbb{C}^2$ fibred over $\mathbb{C}$, the LS manifold associated to $\phi(\theta z)$ is

$$\Lambda_\theta = \{(z, \frac{\partial}{\partial z}\phi(\theta z)) : z \in \mathbb{C}\} = \{(z, \theta\frac{\partial \phi}{\partial z}(\theta z)) : z \in \mathbb{C}\}.$$

Now consider the $\mathbb{C}^*$-action on $\mathbb{C}^2$ given by

$$\rho(\lambda) : (z,w) \mapsto (\lambda z, \lambda^{-1} w), \quad \lambda \in \mathbb{C}^*$$

which also restricts to an action of $S^1 = \partial D$. Thus $\rho(\theta)$ for $\theta \in S^1$ maps Λ_θ to Λ_1.

We will prove below that the HMAE with boundary condition F has a regular solution. Assuming this for now, we show how to produce the desired solution to the Hele-Shaw flow. By Theorem 8.1 there is an associated family of holomorphic discs g_x in $W_{\mathbb{P}^1}$. Since the function $F(z,\theta)$ is invariant under the S^1-action ρ, it implies that the solution Φ is invariant, and in particular the restriction Φ_0 is S^1-invariant. Set $\phi_0 := \Phi_0 + \log(1+|z|^2)$. Then ϕ_0 is an S^1-invariant strictly subharmonic function, which implies that $\frac{\partial \phi_0}{\partial z}(0) = 0$ and $\frac{\partial \phi_0}{\partial z}(z) \neq 0$ whenever $z \neq 0$. This means that $g_0(0) = (0,0)$, while for any $x \neq 0$, the second component of $g_x(0)$ is not 0.

Furthermore the family of discs g_x are be invariant under ρ, i.e.

$$\rho(\theta) g_x(\theta\tau) = g_{\theta x}(\tau) \quad \text{for } \theta \in \partial D, \tau \in D.$$

In particular

$$g_0(\theta) = \rho(\theta^{-1})(g_0)(1) \quad \text{for } \theta \in \partial D \tag{14}$$

and since $\pi(g_0(0)) = 0$ we see πg_0 must be the constant map to 0. So, by continuity of the family g, we have that as long as $|x|$ is sufficiently small, the image of $g_x : D \to W_{\mathbb{P}^1}$, $x \neq 0$, lies in $W_{\mathbb{C}} \cong \mathbb{C}^2$.

So fix such an x with $x \neq 0$. Then by twisting the disc using the $\mathbb{C}^*$-action ρ we get a new punctured disc $f_x : D^\times \to \mathbb{C} \times \mathbb{P}^1$,

$$f_x(\tau) := \rho(\tau)(g_x(\tau)).$$

Since the second coordinate of $g_x(0)$ is not zero f_x extends to a map $D : \mathbb{C} \times \mathbb{P}^1$. Moreover for $\theta \in \partial D$ and $\tau \in D$ we have

$$f_{\theta x}(\tau) = \rho(\tau) g_{\theta x}(\tau) = \rho(\tau\theta) g_x(\theta\tau) = f_x(\theta\tau)$$

so the holomorphic disc defined by f_x depends only on $t = |x|$, which we shall now denote by Σ_t. Observe that $f_x(0) = (0,\infty)$ so Σ_t meets $\{0\} \times \mathbb{P}^1$ at $(0,\infty)$. Moreover if $\tau \in \partial D$ then $g_x(\tau) \in \Lambda_\tau$ so the boundary of Σ_t lies in Λ_1.

Recall that the map $\gamma(x) := \pi g_x(1)$ is a diffeomorphism of $\mathbb{P}^1$ to itself. Since, by invariance $\pi f_x(\theta) = \pi g_{\theta x}(1) = \gamma(\theta x)$ for $\theta \in \partial D$ we deduce

$$\partial\Omega_t = \{\pi f_x(\theta) : \theta \in \partial D\} = \{\gamma(\theta x) : \theta \in \partial D\}$$

which is precisely the image of the circle of radius $t = |x|$ under γ. In particular this shows that, for small $t > 0$, Ω_t is an increasing family of smooth Jordan domains containing the origin, and $\pi\colon \Sigma_t \to \pi(\Sigma_t)$ is an isomorphism and the limit Ω_t as t tends to zero is the origin. Moreover by Theorem 3.10 the family $\Omega_t = \pi(\Sigma_t)$ satisfies the Hele-Shaw flow as required, after possible reparameterization of t.

Thus it remains to prove that there is regular solution to the HMAE with boundary data F in (13). Observe that if ϕ happens to be rotation invariant, i.e. $\phi(\tau z) = \phi(z)$ for all $\tau \in \partial D$, the HMAE has a trivial regular solution, namely $\Phi(z,\tau) := \phi(z) - \log(1+|z|^2)$. (This reflects the trivial observation that if the permeability κ is radially symmetric then the Hele-Shaw flow with empty initial condition will be the trivial solution in which Ω_t is a disc centered at the origin.)

Let ψ denote a rotation invariant potential that we also assume to be equal to $|z|^2$ in a neighbourhood of the origin. By Donaldson's theorem we know that there is a C^2-ball of radius $\epsilon > 0$ centred at $\phi_0 - \log(1+|z|^2)$ consisting of boundary conditions for which the HMAE has a regular solution.

Now an arbitrary smooth strictly subharmonic potential ϕ can be written as

$$\phi(z) = \mathrm{Re}(a + bz + c\bar{z} + dz^2 + e\bar{z}^2) + f|z|^2 + o(|z|^2),$$

where f is some real positive number. Our requirement on ϕ was that $\Delta\phi = 1/\kappa$, so without loss of generality we can assume that this first term is zero since it is harmonic, and also by rescaling that $f = 1$. This means that on some small disc centered at the origin ϕ lies in the C^2-ball of radius $\epsilon/2$ around ψ.

Pick δ_1 and δ_2 to be small positive numbers so that ϕ is $\epsilon/2$ close to $(1+\delta_1)\psi - \delta_2$, and so that on the boundary of the small disc $\phi < (1+\delta_1)\psi - \delta_2$. Still, in a neighbourhood of 0, we will have that $\phi > (1+\delta_1)\psi - \delta_2$. Let f_δ be a smooth non-negative function on the real line of integral one and with compact support in a δ-neighbourhood of 0. Then let ϕ_δ denote the subharmonic function

$$\phi_\delta := \int_\lambda \max(\phi, (1+\delta_1)\psi - \delta_2 + \lambda) f(\lambda) d\lambda.$$

This new potential will be smooth and strictly subharmonic. If δ is chosen small enough, $\phi_\delta = \phi$ in a neighbourhood of the origin, while in a neighbourhood of the boundary of the disc it will be equal to $(1+\delta_1)\psi + c$ where c is some constant (the integral of $\lambda f \lambda$ minus δ_2). We can thus extend it by letting it be equal to $(1+\delta_1)\psi + c$ outside of the disc. For sufficiently small δ, $\phi_\delta - (1+\delta_1)\psi - c$ will have C^2-norm of less than ϵ, and thus the same is true for $\tilde{F} := \frac{\phi_\delta}{(1+\delta_1)} - \log(1+|z|^2)$.

This implies that the HMAE with $\tilde{F}$ as boundary condition has a regular solution Φ. By scaling, letting $(1+\delta_1)\omega_0$ be the new background Kähler form, the HMAE with boundary condition $F = (1+\delta_1)F$ will have $(1+\delta_1)\Phi$ as a regular solution as well, completing the proof.

References

[1] D. Aharanov and H. S. Shapiro *Domains in which analytic functions satisfy quadrature identities* J. Analyse Math. 30 (1976), 39-73.

[2] S. N. Antontsev, C. R. Gonçalves and A. M. Meirmanov *Local existence of classical solutions to the well-posed Hele-Shaw problem* Port. Math. (N.S.) 59 (2002), no. 4, 435-452.

[3] E. Bedford and M. Kalka *Foliations and complex Monge-Ampére equations* Comm. Pure Appl. Math., 30 (1977) no 5, 543-571.

[4] R Berman *Bergman kernels and equilibrium measures for line bundles over projective manifolds* Amer. J. Math. 131 (2009), no. 5, 1485–1524.

[5] E. Bishop *Differentiable manifolds in complex Euclidean space* Duke Math. Jour., 32 (1965), 1-22.

[6] V. G. Cherednichenko *Inverse Logarithmic Potential Problem*. Walter de Gruyter (1996) ISBN 9067642029.

[7] S. Cho and S. R. Pai *On the regularity of the Riemann mapping function in the plane* Pusan-Kyongnam Math. J. 12 (1996), 203-211.

[8] P. Davis *The Schwarz function and its applications* Carus Math. Monographs 17. Math. Assoc. of America, 1974.

[9] S. K. Donaldson *Holomorphic discs and the complex Monge-Ampère equation* J. Symplectic Geom., 1 (2002), no. 2, 171-196.

[10] C. M. Elliott and V. Janovsky *A variational inequality approach to Hele-Shaw flow with a moving boundary* Proc. Roy. Soc. Edin., A88 (1981), 93-107.

[11] J. Escher and G. Simonett *On Hele-Shaw models with surface tension* Math. Res. Lett., 3 (1996), no. 4, 467-474.

[12] J. Escher and G. Simonett *Classical solutions for Hele-Shaw models with surface tension* Adv. Differential Equations, 2 (1997), no. 4, 619-642.

[13] J. Escher and G. Simonett *Classical solutions of multidimensional Hele-Shaw models* SIAM J. Math. Anal. 28 (1997), no. 5, 1028-1047.

[14] F. Forstnerič *Analytic disks with boundaries in a maximal real submanifold of* $\mathbb{C}^2$ Ann. Inst. Fourier (Grenoble), 37 (1987), no. 1, 1-44.

[15] F. D. Gakhov *Boundary value problems* Pergamon Press, Oxford-New York-Paris, Addison-Wesley Publishing Co., Inc., Reading, Mass.-London 1966.

[16] M. Gromov *Pseudoholomorphic curves in symplectic manifolds* Invent. Math., 82 (1985), 307-347.

[17] B. Gustafsson *Quadrature identities and the Schottky double* Acta Appl. Math., 1 (1983), no. 3, 209-240.

[18] B. Gustafsson *On a differential equation arising in a Hele-Shaw flow moving boundary problem* Arkiv för Mat., 22 (1984), no. 1, 251-268.

[19] B. Gustafsson *Applications of variational inequalities to a moving boundary problem for Hele-Shaw flows* SIAM J. Math. Analysis, 16 (1985), no. 2, 279-300.

[20] B. Gustafsson and A. Vasil'ev *Conformal and potential analysis in Hele-Shaw cells* Advances in Mathematical Fluid Fluid Mechanics. Birkhäuser Verlag, Basel, 2006.

[21] E. Hanzawa *Classical solutions of the Stefan problem* Tohoku Math. J. (2), 33 (1981), no. 3, 297-335.

[22] H. Hedenmalm and S. Shimorin *Hele-Shaw flow on hyperbolic surfaces* J. Math. Pures Appl. (9) 81 (2002), no. 3, 187-222.

[23] H. Hedenmalm and A. Olofsson *Hele-Shaw flow on weakly hyperbolic surfaces* Indiana Univ. Math. J. 54 (2005), no. 4, 1161-1180.

[24] V. Isakov *Inverse Source Problems* Mathematical Surveys and Monographs. Number 34. American Mathematical Society, 1990. ISBN: 0-8218-1532-6

[25] D. Khavinson, M. Mineev-Weinstein and M Putinar *Planar elliptic growth* Complex Anal. Oper. Theory 3 (2009), no. 2, 425-451.

[26] C. LeBrun and L. J. Mason *Zoll manifolds and complex surfaces* J. Differential Geom., 61 (2002), no. 3, 453-535.

[27] C. LeBrun *Twistors, holomorphic discs, and Riemann surfaces with boundary* Perspectives in Riemannian geometry, 209-221, CRM Proc. Lecture Notes, 40, Amer. Math. Soc., Providence, RI, 2006.

[28] Y.-L. Lin *Perturbation theorems for Hele-Shaw flows and their applications* Arkiv för Mat., 49 (2011), no. 2, 357-382.

[29] D. McDuff and D. Salamon *J-holomorphic curves and Symplectic Topology* American Mathematical Society Colloquium Publications, 52. Americal Mathematical Society, Providence, RI, 2004.

[30] M. Reissig and L. von Wolfersdorf *A simplified proof for a moving boundary problem for Hele-Shaw flows in the plane* Ark. Mat. 31 (1993), 101-116.

[31] S. Richardson *Hele-Shaw flows with a free boundary produced by the injection of fluid into a narrow channel* J. Fluid Mech., 56 (1972), no. 4, 609-618.

[32] M. Sakai *Quadrature domains* Lecture notes in Math. 934, Springer-Verlag, New York, 1982.

[33] M. Sakai *Regularity of boundaries having a Schwarz function* Acta Math. 166 (1991), 263-297.

[34] M. Sakai *Regularities of boundaries in two dimensions* Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 20 (1993), 323-339.

[35] M. Sakai *Application of variational inequalities to the existence theorem on quadrature domains* Trans. Amer. Math. Soc. 276 (1993), 267-279.

[36] M. Sakai *Regularity of boundaries of quadrature domains in two dimensions* SIAM J. Math. Analysis 24 (1993), no. 2, 341-364.

[37] H. S. Shapiro *The Schwarz function and its generalization to higher dimensions* University of Arkansas Lecture Notes in the Mathematical Sciences, 9. A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York, 1992.

[38] F. R. Tian *A Cauchy integral approach to Hele-Shaw problems with a free boundary* Arch. Rational Mech. ANal. 135 (1996), no. 2, 175-196.

[39] F. R. Tian *Hele-Shaw problems in multidimensional spaces* J. Nonlinear Science 10 (2000), 275-290.

[40] Y. P. Vinogradov and P. P. Kufarev *On a problem of filtration* Akad. Nauk SSSR. Prikl. Mat. Meh., 12 (1948), 181-198. (in Russian)

[41] P. B. Wiegmann and A. Zabrodin *Conformal maps and dispersonless integrable hierarchies* Preprint (1999) arXiv:hep-th/9909147.

JULIUS ROSS, UNIVERSITY OF CAMBRIDGE, UK.
J.ROSS@DPMMS.CAM.AC.UK

DAVID WITT NYSTROM, UNIVERSITY OF GOTHENBURG, SWEDEN.
WITTNYST@CHALMERS.SE